



Evaluation of Organic Inputs Viz., Jeevamruth and Bijamruth on Soil Nutrients and Uptake by Fenugreek Absorption

Sugumaran M.P ^{a*}, Goveanthan A.S ^b, Bharathi Kumar K ^{c*},
Kiruba M ^{b*}, Akila S ^a, Somasundaram E ^a, Natarajan K ^a,
Gayathry G ^a and Sinduja M ^d

^a Krishi Vigyan Kendra, TNAU, Vridhachalam, India.
^b Tamil Nadu Agricultural University, Coimbatore, India.
^c Krishi Vigyan Kendra, TNAU, Sandhiyur, India.
^d National Agro Foundation, Chennai, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.56557/ajocr/2025/v10i19043>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.ikpress.org/review-history/12646>

Original Research Article

Received: 04/11/2024
Accepted: 08/01/2025
Published: 13/01/2025

ABSTRACT

The present study on the 'Biochemical evaluation of organic inputs (Jeevamruth and Beejamruth) and their efficacy on Greens' was conducted at the Department of Sustainable Organic Agriculture, Tamil Nadu Agricultural University, Coimbatore, with laboratory experiments carried out at the

*Corresponding author: E-mail: sugumaran.mp@tnau.ac.in, kiruba.m@tnau.ac.in, bharathisolunum@gmail.com;

Cite as: M.P, Sugumaran, Goveanthan A.S, Bharathi Kumar K, Kiruba M, Akila S, Somasundaram E, Natarajan K, Gayathry G, and Sinduja M. 2025. "Evaluation of Organic Inputs Viz., Jeevamruth and Bijamruth on Soil Nutrients and Uptake by Fenugreek Absorption". *Asian Journal of Current Research* 10 (1):15-21. <https://doi.org/10.56557/ajocr/2025/v10i19043>.

Department of Environmental Science. A field trial was set up in a randomized block design with three replications. Biometric observations were recorded on the 10th, 20th, and 30th days after sowing. Soil samples were collected initially and at harvest, and analyzed for chemical and biological properties. Plant samples were tested for protein and chlorophyll content. The application of Jeevamruth increased soil nutrients (NPK) and organic carbon levels. Liquid organic preparations were found to have higher populations of bacteria, fungi, and actinomycetes. The findings suggest that Beejamruth should be applied on the day of preparation, whereas Jeevamruth should be used within 10 to 15 days of preparation. The use of these liquid formulations helps supplement nutrients when biofertilizers are applied.

Keywords: Jeevamruth; Beejamruth; biochemical evaluation; organic inputs; organic farming; green revolution.

1. INTRODUCTION

“The Green Revolution intensified agriculture to meet the growing demand for food and fiber, but this practice has come at a significant environmental cost, leading to the ongoing loss of natural ecosystems, depletion of groundwater, food contamination, and other forms of environmental degradation. A major challenge for the nation in the coming years will be to provide safe food for the expanding population. In this context, organic farming, a holistic production management system designed to enhance the health of agroecosystems, has gained widespread recognition as a viable alternative to conventional food production and ensures safe food for human consumption. This farming approach minimizes the use of synthetic fertilizers, growth regulators, and livestock feed additives, and instead relies on green manures, crop rotations, crop residues, animal manures, biofertilizers, and various cow-based liquid organic manures such as Panchagavya, Jeevamruth, Beejamruth, Amritpani, etc. Among the liquid formulations, Panchagavya is one of the most significant to consider for shelf life studies, as it is not only highly effective in promoting crop growth but also widely utilized by farmers”. (Sugumaran et al., 2018). “The compounds Erioflorin and Nagilactone A present in Beejamruth act as plant growth regulators, making Beejamruth a valid and effective alternative fertilizer for producing safe, high-quality food that meets the needs of modern Indian agriculture”. (Goveanthan et al., 2019). “Compounds such as Isoenanthic acid, Columbianetin, Lomatin, 1,6-Hexanediol, Mevastatin, Gitoxigenin, Dibutoxy anthracene, Erioflorin, Nagilactone, Trimegestone, Rofe Coxib, and Clupanodonic acid present in Jeevamruth aid in plant metabolism and significantly enhance its growth”. (Goveanthan et al., 2021).

“Organic agriculture is increasingly gaining recognition as a key component of development, demonstrating considerable promise commercially, socially, and environmentally. While the concept has evolved over time, the modern organic movement is fundamentally different from its original form. Liquid formulations used in organic agriculture, such as Panchagavya, Beejamruth, and Jeevamruth, are fermented products made from locally available materials that act as plant growth enhancers. These formulations are rich in beneficial microorganisms that support and stimulate plant growth, leading to improved vegetative development and higher-quality yields. Formulations derived from agricultural by-products like grain bran, oil cakes, and farmyard manure serve as effective growth carriers and storage media” (Deva kumar et al., 2011). In recent years, there has been a growing interest in the use of Panchagavya, Beejamruth, Jeevamruth, and other liquid organic formulations in organic farming.

2. MATERIALS AND METHODS

2.1 Field Experimental Details

The experiment was carried out using a randomized block design with three replications. The experimental layout remained undisturbed throughout the investigation period, and fenugreek seeds were soaked for 1 hour in the Beejamruth solution prior to sowing, after which they were planted in the field.

Design	: RBD
Number of treatments	: 7
Number of replications	: 3

The treatment details are given;

2.2 Treatments

- T₁–Control
- T₂–Jeevamruth @ 3% Spray (Green gram flour)
- T₃–Jeevamruth @ 5 % Spray (Green gram flour)
- T₄–Jeevamruth @ 3 % Spray (Black gram flour)
- T₅–Jeevamruth @ 5 % Spray (Black gram flour)
- T₆–Jeevamruth @ 3 % Spray (Green gram + Black gram flour)
- T₇–Jeevamruth @ 5% Spray (Green gram +Black gram flour)

2.3 Preparation of Soil Sample

Composite soil samples were collected before the experiment and analyzed for their mechanical and chemical properties. After harvest, soil samples were collected from each treatment plot and analyzed for pH, EC, organic carbon, and major nutrients following standard procedures.

2.4 Preparation of the Plant Sample

The samples were collected to estimate dry matter production, which was used for calculating nutrient uptake. Oven-dried plant samples were ground using a Wiley-Mill, sieved, and analyzed to determine the total NPK uptake by multiplying the N, P, and K contents with the dry matter at each respective stage. The uptake values were then calculated and reported. The samples were pulverized and sieved through a 0.2 mm mesh sieve, and the analyses were performed following standard procedures.

3. RESULTS AND DISCUSSION

The results from the experiment on “Biochemical evaluation of organic inputs (Jeevamruth and Beejamruth) and their efficacy on greens” conducted at the Department of Sustainable Organic Farming, Tamil Nadu Agricultural University, Coimbatore are summarized.

3.1 Soil Parameters

The soil pH was not significantly influenced by the different treatments. However, numerically higher soil pH was observed as given in the (Table 1). The Jeevamruth, as organic source of nutrient slightly increased the soil pH. But no significant difference in pH among various treatments was noticed. However, treatment which received Jeevamruth @ 5 % Spray (Green gram flour) recorded higher pH among the treatments. This was supported by Elias-Azar,

(1980) who reported that soil pH had increased due to application of poultry manure as organic nutrient source. The soil EC was not significantly influenced by the different treatments. The high / low EC value was found in Jeevamruth @ 5 % Spray (Green gram flour) and Jeevamruth @ 5 % Spray (Black gram flour). At the time of inception of the study, the soil organic content was 0.28 per cent. After harvest of the fenugreek crop, the organic carbon content of soil was higher in Jeevamruth @ 5% Spray (Green gram flour) (1.08%), followed by Jeevamruth @ 5 % Spray (Green gram +Black gram flour) (0.92%). The lowest organic carbon content was observed in control (0.33%). Addition of Jeevamruth @ 5 % Spray (Green gram flour) resulted in significant increase in the organic carbon content of soil. These results are in agreements with the findings of Mathan, (2000) and Maskina et al., (1988).

The available soil nitrogen was not significantly influenced by the sources of nutrition. Numerically higher soil nitrogen was recorded in the treatments Jeevamruth @ 3% Spray (Green gram flour) and Jeevamruth @ 5 % Spray (Green gram +Black gram flour). The highest soil available phosphorus at 30 DAS, was recorded in the Jeevamruth @ 5% Spray (Green gram flour) and Jeevamruth @ 5 % Spray (Green gram +Black gram flour) treatments with values of 19.0 and 16.0 kg ha⁻¹ respectively. The high soil potassium content was recorded in Jeevamruth @ 5% Spray (Green gram flour) (264.0 kg ha⁻¹) followed by Jeevamruth @ 5 % Spray (Green gram +Black gram flour) (239.0 kg ha⁻¹) treatments. Sole cropping with biofertilizers produced the highest seed yields for fennel (2233 kg ha⁻¹) and fenugreek (1240 kg ha⁻¹) (Ghaderimokri et al., 2022).

3.2 Plant Total NPK

The analysis for total NPK in fenugreek after harvest showed that the total nitrogen content was found to be high in Jeevamruth @ 5% Spray (Green gram +Black gram flour) with 1.5 per cent, whereas the total phosphorus content was high in Jeevamruth @ 5% Spray (Green gram) (0.33 per cent). The potassium content was high in the treatments Jeevamruth @ 5% Spray (Green gram), Jeevamruth @ 5% Spray (Black gram flour) and Jeevamruth @ 5% Spray (Green gram +Black gram flour) with 0.22 per cent and they were on par with each other (Table 2). “Nutrient uptake is a cordially event of nutrient concentration and dry matter accumulation.

Organic manures promoted nutrient utilization and accounts for better NPK uptake. Increased uptake might be due to higher availability of nutrients from the soil reservoir and also from the added sources of organic manures” (Priyadarsini & Prasad, 2003). The rate of uptake is dependent upon by crop N demand, phenological stage, soil N availability, transpiration, rooting depth and soil water status. Crop nitrogen demand is estimated depending on the rate of growth and the maximum concentration of nitrogen that different organic can accumulate depends upon their composition. Nitrogen uptake was maximum in Jeevamruth @ 5 % Spray (Green gram) and it was comparable with Jeevamruth @ 5 % Spray (Green gram +Black gram flour). Organic inputs known to have a favorable effect on soil structure, texture and tilth thus facilitate quick and greater availability of plant nutrients and provides a better environment for root growth and proliferation, thereby creating more absorptive surface for uptake of nutrients. These results are in conformity with the findings of Chavan et al., (1997); Shashidhara, (2000) and Kuttimani, (2004) in chillies. The organic inputs might have increased the soil organic P content leading to increased P availability. Higher phosphorus uptake was recorded in T₃ which was on par with T₇. Increased P availability might be due to solubilisation of native P by the organic acids produced during organic inputs decomposition, thus leading to better utilization of available P, which in turn favored better P uptake. Similar results were also obtained by Beulah, (2001), Sreekhantan, (1987) and Somasundaram, (1991). The highest K uptake was registered with Jeevamruth @ 5 % spray (Use Green gram flour for preparation). The lower level of K uptake was observed in control (no manure/ no spray) at all the growth stages of

the crop. The increased uptake of K observed in above said treatments might be the result of increased availability of K in soils due to the basal application of enriched farm yard manure. The enhanced K availability irrespective of the season coupled with higher K uptake due to organic manure incorporation could be attributed to higher DMP and K absorption, evidencing the priming effect of K contribution by organic manure. These results are in conformity with the findings of Santos & Nicoloso, (1990) and Kuttimani, (2004). “The plant height, root length and single plant weight in fenugreek are high in the treatment as Jeevamruth 5% spray was observed as a viable organic approach to improve soil and eco-friendly fenugreek production” (Goveanthan et al., 2020). “In another study, seeds treated with Panchagavya and Jeevamruth separately, the maximum shoot length and root length were recorded in Panchagavya treatment and minimum shoot length and root length of was recorded in Jeevamruth treated seeds and also the panchagavya treated seeds registered the maximum vigour index” (Akila et al., 2020). “In another study, the soluble protein content (0.87 mg/g) and total sugar content (11.20 µg/g) were found to be improved in Panchagavya (Groundnut cake instead of ghee) @ 3% spray treatment and Recommended dose of NPK fertilizer applied plants” (Sugumaran et al., 2019). “Another study showed that a 2:1 fenugreek-buckwheat intercropped system with the application of integrated fertilizer and broiler litter can successfully be implemented for improving productivity, N and P contents of fenugreek and buckwheat as well as the nutrient land equivalent ratio (compared with sole cropping with chemical fertilizer) under semi-arid growing conditions” (Salehi et al., 2018).

Table 1. Effect of organic nutrient source (Jeevamruth) on soil pH, EC (dS m⁻¹) and organic carbon (%)

Treatment	pH	EC (dSm ⁻¹)	Organic Carbon (%)
T ₁ - Control	7.26	0.3	0.33
T ₂ - Jeevamruth @ 3% Spray (Green gram)	7.28	0.8	1.08
T ₃ - Jeevamruth @ 5 % Spray (Green gram)	7.29	1.5	0.90
T ₄ - Jeevamruth @ 3 % Spray (Black gram)	7.40	0.4	0.46
T ₅ - Jeevamruth @ 5 % Spray (Black gram)	7.62	1.5	0.72
T ₆ - Jeevamruth @ 3 % Spray (Green gram + Black gram)	7.58	0.4	0.52
T ₇ - Jeevamruth @ 5 % Spray (Green gram + Black gram)	7.74	0.8	0.92
SEd	0.0021	0.0055	0.003
CD (P = 0.05)	0.0045	0.0119	0.06

Table 2. Effect of organic nutrient source (Jeevamruth) on fenugreek plant total nitrogen, total phosphorus and total

Treatment	Nitrogen (%)	Phosphorus (%)	Potassium (%)
T ₁ - Control	0.16	0.05	0.17
T ₂ - Jeevamruth @ 3% Spray (Green gram)	0.70	0.12	0.20
T ₃ - Jeevamruth @ 5 % Spray (Green gram)	1.23	0.33	0.23
T ₄ - Jeevamruth @ 3 % Spray (Black gram)	1.00	0.16	0.20
T ₅ - Jeevamruth @ 5 % Spray (Black gram)	1.50	0.25	0.22
T ₆ - Jeevamruth @ 3 % Spray (Green gram + Black gram)	1.06	0.18	0.21
T ₇ - Jeevamruth @ 5 % Spray (Green gram + Black gram)	1.50	0.31	0.22
SEd	0.0051	0.0011	0.0002
CD (P = 0.05)	0.0112	0.0024	0.0005




Fig. 1. Effect of organic nutrient source (Jeevamruth) on soil available nitrogen, available phosphorus and available potassium (kg ha⁻¹)



Fig. 2. Experimental field view

4. CONCLUSIONS

The field experiment indicates that plant height, root length, and single plant weight were highest in the T3 treatment (Jeevamruth @ 5% spray). Applying Jeevamruth to the soil enhanced nutrient (NPK) levels and organic carbon content. Liquid organic formulations are rich in bacteria, fungi, and actinomycetes. Studies suggest Beejamruth should be applied on the day of preparation, while Jeevamruth remains effective for 10 to 15 days after preparation. These liquid formulations can effectively supplement nutrients when combined with biofertilizers.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

1. ChatGpt was used only for editing this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Akila, S., Sugumaran, M. P., Suganya, K., & Somasundaram, E. (2020). Studies on testing the efficacy of liquid organic inputs (Panchagavya and Jeevamruth) on maize (*Zea mays* L.) germination. *Current Journal of Applied Science and Technology*, 39(23), 134-137.
- Beulah, A. (2001). *Growth and development of moringa (Moringa oleifera Lem.) under organic and inorganic system of culture* (Ph.D. thesis). Tamil Nadu Agricultural University, Coimbatore.
- Chavan, P., Syedismaiz, J., Rudraksha, G. B., Malewar, G., & Baig, M. I. (1997). Effect of various nitrogen levels through FYM and urea on yield, uptake of nutrients, and ascorbic acid content in chilli (*Capsicum annuum* L.). *Journal of the Indian Society of Soil Science*, 45, 833-835.
- Deva Kumar, N., Rao, G. G. E., & Shuba, S. (2011). Evaluation of locally available media for the growth and development of nitrogen fixing microorganisms. In *Proceedings of the 3rd scientific conference of ISOFAR Organic are life knowledge for tomorrow* (pp. 504-509). Korea.
- Elias-Azar, K. (1980). Biocarbonate extractable phosphorus in fresh and composted dairy manure. *Soil Science Society of America Journal*, 44(2), 434-435.
- Ghaderimokri, L., Rezaei-Chiyaneh, E., Ghiyasi, M., Gheshlaghi, M., Battaglia, M. L., & Siddique, K. H. (2022). Application of humic acid and biofertilizers changes oil and phenolic compounds of fennel and fenugreek in intercropping systems. *Scientific Reports*, 12(1), 5946. <https://doi.org/10.1038/s41598-022-10111-w>
- Goveanthan, A. S., Sugumaran, M. P., & Somasundaram, E. (2019). Biochemical analysis of Beejamruth and its plant promoting factors. *International Journal of Current Research and Academic Review*, 7(5), 1-4.
- Goveanthan, A. S., Sugumaran, M. P., & Somasundaram, E. (2021). Scientific validation of organic liquid formulation-Jeevamruth by studying its characteristics. *International Journal of Plant Sciences*, 16(1), 15-18.
- Goveanthan, A. S., Sugumaran, M. P., Gudimetha, G. K., Akila, S., Suganya, K., & Somasundaram, E. (2020). Studies on organic inputs (Jeevamruth and Beejamruth) and their efficacy on fenugreek. *The Pharma Innovation Journal*, 9(11), 92-94.
- Kuttimani, S. (2004). *Response of chilli (Capsicum annuum L.) genotypes to integrated nutrient management* (M.Sc. thesis). University of Agricultural Sciences, Dharwad.
- Maskina, M. S., Singh, Y., & Singh, B. (1988). Response of wetland rice to fertilizer N in soil amended with cattle, poultry, and pig manure. *Bio-wastage*, 26(1), 1-8.
- Mathan, K. K., Appavu, K., & Saravanan, A. (2000). Effect of organics and irrigation levels on soil physical properties and yield of crops under sorghum-soybean cropping system. *Madras Agricultural Journal*, 87(1-3), 50-53.
- Priyadarsini, J., & Prasad, P. V. N. (2003). Evaluation of nitrogen use efficiency of

- different rice varieties supplied with organic and inorganic sources of nitrogen. *Andhra Agricultural Journal*, 50(4), 207-210.
- Salehi, A., Mehdi, B., Fallah, S., Kaul, H. P., & Neugschwandtner, R. W. (2018). Productivity and nutrient use efficiency with integrated fertilization of buckwheat–fenugreek intercrops. *Nutrient Cycling in Agroecosystems*, 110, 407-425. <https://doi.org/10.1007/s10705-018-9922-3>
- Santos, O. S. Dos, & Nicoloso, F. T. (1990). Effects of mineral nitrogen, molybdenum, and inoculation with rhizobium on common beans. *Revista do Centro de Ciências Rurais Universidade Federal de Santa Maria*, 20(1-2), 23-25.
- Shashidhara, G. B. (2000). *Integrated nutrient management for chilli (Capsicum annum L.) in alfisols of the Northern transition zone of Karnataka* (M.Sc. thesis). University of Agricultural Sciences, Dharwad.
- Somasundaram, E. (1991). Studies on the direct and residual effect of applied and intercropped *Sesbania rostrata* on rice. *Asian Journal of Plant Sciences*, 6(2), 282-287.
- Sreekhantan, L. (1987). *Integrated phosphorus management in rice-based cropping system* (Ph.D. thesis). Tamil Nadu Agricultural University, Coimbatore.
- Sugumaran, M. P., Akila, S., & Somasundaram, E. (2018). Studies on analyzing the shelf life of Panchagavya with different alternatives for ghee. *International Journal of Agriculture Sciences*, 10(24), 7655-7656.
- Sugumaran, M. P., Akila, S., & Somasundaram, E. (2019). Studies on analysis of biochemical characters of leaf over liquid organic inputs (Panchagavya and Jeevamruth) on maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*, 8(5), 1794-1797.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://prh.ikpress.org/review-history/12646>